

# Phylogenomics Revives Traditional Views on Deep Animal Relationships

Hervé Philippe,<sup>1,11</sup> Romain Derelle,<sup>2,11</sup> Philippe Lopez,<sup>2</sup> Kerstin Pick,<sup>3,5</sup> Carole Borchiellini,<sup>6</sup> Nicole Boury-Esnault,<sup>6</sup> Jean Vacelet,<sup>6</sup> Emmanuelle Renard,<sup>6</sup> Evelyn Houliston,<sup>7</sup> Eric Quéinnec,<sup>2</sup> Corinne Da Silva,<sup>8</sup> Patrick Wincker,<sup>8</sup> Hervé Le Guyader,<sup>2</sup> Sally Leys,<sup>9</sup> Daniel J. Jackson,<sup>3,10</sup> Fabian Schreiber,<sup>4</sup> Dirk Erpenbeck,<sup>5</sup> Burkhard Morgenstern,<sup>3,4</sup> Gert Wörheide,<sup>5,\*</sup> and Michaël Manuel<sup>2,\*</sup>

<sup>1</sup>Centre Robert-Cedergren  
Département de Biochimie  
Université de Montréal  
Succursale Centre-Ville  
Montréal, Québec H3C3J7  
Canada

<sup>2</sup>UPMC, Univ Paris 06  
UMR 7138 Systématique, Adaptation, Evolution  
CNRS UPMC MNHN IRD, Case 05  
Université Pierre et Marie Curie  
7 quai St Bernard  
75005 Paris  
France

<sup>3</sup>Courant Research Center Geobiology  
Georg-August-Universität Göttingen Goldschmidtstr. 3  
<sup>4</sup>Abteilung Bioinformatik  
Institut für Mikrobiologie und Genetik  
Goldschmidtstr. 1  
37077 Göttingen  
Germany

<sup>5</sup>Department of Earth- and Environmental Sciences & GeoBioCenter<sup>LMU</sup>  
Ludwig-Maximilians-Universität München  
Richard-Wagner-Str. 10  
80333 München  
Germany

<sup>6</sup>Aix-Marseille Université  
CNRS UMR 6540 DIMAR  
Centre d'Océanologie de Marseille  
Station Marine d'Endoume  
rue de la Batterie des Lions  
13 007 Marseille  
France

<sup>7</sup>UPMC, Univ Paris 06  
CNRS UMR 7009 Biologie du Développement  
Observatoire Océanologique  
06230 Villefranche-sur-Mer  
France

<sup>8</sup>Genoscope and CNRS UMR 8030  
2 rue Gaston Crémieux  
91057 Evry  
France

<sup>9</sup>Department of Biological Sciences  
CW 405  
University of Alberta

Edmonton, AB T6G 2E9  
Canada

<sup>10</sup>School of Integrative Biology  
The University of Queensland  
Brisbane 4072  
Australia

## Summary

The origin of many of the defining features of animal body plans, such as symmetry, nervous system, and the mesoderm, remains shrouded in mystery because of major uncertainty regarding the emergence order of the early branching taxa: the sponge groups, ctenophores, placozoans, cnidarians, and bilaterians. The “phylogenomic” approach [1] has recently provided a robust picture for intrabilaterian relationships [2, 3] but not yet for more early branching metazoan clades. We have assembled a comprehensive 128 gene data set including newly generated sequence data from ctenophores, cnidarians, and all four main sponge groups. The resulting phylogeny yields two significant conclusions reviving old views that have been challenged in the molecular era: (1) that the sponges (Porifera) are monophyletic and not paraphyletic as repeatedly proposed [4–9], thus undermining the idea that ancestral metazoans had a sponge-like body plan; (2) that the most likely position for the ctenophores is together with the cnidarians in a “coelenterate” clade. The Porifera and the Placozoa branch basally with respect to a moderately supported “eumetazoan” clade containing the three taxa with nervous system and muscle cells (Cnidaria, Ctenophora, and Bilateria). This new phylogeny provides a stimulating framework for exploring the important changes that shaped the body plans of the early diverging phyla.

## Results and Discussion

### A Comprehensive Phylogenomic Data Set to Address Basal Metazoan Evolution

Previous studies of basal metazoan relationships by molecular phylogeny techniques (e.g., [3–8, 10, 11]) have proposed contradictory and often poorly supported trees, leaving major issues such as the phylogenetic status (monophyly or paraphyly) of sponges and the position of ctenophores and placozoans unsettled. These inconsistencies may reflect insufficient molecular sampling and/or inadequate taxon sampling of the diversity of extant nonbilaterian metazoan lineages [1, 11–13]. We have adopted a phylogenomic approach specifically aimed at clarifying the basal metazoan relationships, involving more comprehensive sampling of all the major early branching animal lineages. By using newly generated cDNA sequences in addition to publicly available sequences, we have assembled a metazoan data set enriched in species representing the early diverging phyla (see [Experimental Procedures](#) and [Supplemental Data](#) available online). The data set comprises 128 different protein-coding genes (30,257 unambiguously aligned

\*Correspondence: woerheide@lmu.de (G.W.), michael.manuel@snv.jussieu.fr (M.M.)

<sup>11</sup>These authors contributed equally to this work




Figure 1. Phylogenetic Analyses of 128 Nuclear-Encoded Proteins

Bayesian tree obtained from the analysis of 30,257 aligned amino acid positions for the 55 terminal taxa with the CAT model. Bootstrap supports (BS) after 100 replicates are indicated for three analyses with different taxon sampling: outgroup 1 (BS values in pink); outgroup 2 (BS values in blue); unrooted analysis (BS values in black). Nodes with maximal support values in all analyses are indicated by an asterisk. The tree obtained with outgroup 1 is shown here (and in Figure S1 with branch posterior probabilities, PP), whereas trees obtained with outgroup 2 and without outgroup are shown in Figures S2 and S3, respectively. Scale bar indicates number of changes per site.

positions) for 11 outgroup species and 44 metazoans, including 9 sponge species, 3 ctenophores, 9 cnidarians, the placozoan *Trichoplax*, and a representative sampling of bilaterian species. Among the 55 terminal taxa, 24 are complete or nearly complete ( $\leq 5\%$  of missing data), and only 27% of positions in the final alignment are absent (see Table S2). This is the first phylogenomic data set to include all four main sponge lineages: Demospongiae, by far the most species-rich sponge group, is represented by four species, chosen to maximize morphological and phylogenetic diversity; Hexactinellida and Calcispongiae are each represented by two species; and Homoscleromorpha is represented by a chimerical operational taxonomic unit created from two species of the genus *Oscarella*.

### The Sponges Restored as a Monophyletic Group

Our data set was analyzed by Bayesian inference analysis, via the CAT model of sequence evolution [14], conceived to reduce artifacts resulting from mutational saturation and unequal rates of substitution, which are major problems when analyzing ancient events [13, 15]. To explore the effect of outgroup taxa on the metazoan interrelationships obtained, we performed three analyses with different taxon samplings (Figure 1): rooted analysis with a paraphyletic outgroup comprised of the fungi, ichthyosporeans, *Capsaspora*, and choanoflagellates (“outgroup 1;” tree shown in Figure 1 and Figure S1; with bootstrap supports [BS] in pink in Figure 1); analysis rooted with just choanoflagellates, the metazoan sister group [16] (“outgroup 2;” BS in blue in Figure 1, tree in




Figure 2. Characters of the Sponge Body Plan and Their Evolution

(A) Schematic section of an adult sponge (bottom) and SEM picture showing a choanocyte, the sponge collar cell (top, choanocyte from *Chelonaplysilla noevus*, Demospongiae). The arrows indicate the direction of circulation of water in the aquiferous system of the sponge. Abbreviations: atr, atrial cavity; cb, cell body; cc, choanocyte chamber; col, collar of microvilli; ex, exhalant canal; fl, flagellum; in, inhalant canal; mes, mesohyl; osc, osculum (or exhalant orifice); ost, ostium (or inhalant orifice); pin, pinacoderm (thin epithelial layer, limiting the sponge body on its external surface and within the canals); sp, spicule.

(B) Most parsimonious scenario for the evolution of sponge body plan characters, imposed on a scheme of sponge paraphyly.

(C) Most parsimonious scenario assuming sponge monophyly.

In (B) and (C), the gray branches indicate the presence of sponge body plan characters (aquiferous system, internalized choanocyte chambers, pinacoderm) and the black branches indicate the absence of these characters. The gray horizontal line indicates character acquisition; the hollow horizontal line indicates character loss. "Sponges 1, 2, and 3" correspond to the major lineages (silicisponges, homoscleromorphs, and calcisponges), of which exact branching order varies among published studies recovering sponge paraphyly.

Figure S2); and unrooted analysis (BS in black in Figure 1, tree in Figure S3). The topology resulting from the rooted analyses (trees shown in Figure 1 and Figures S1 and S2) was statistically well supported at most nodes, and its general features were in line with previous studies [2, 3]: choanoflagellates positioned as the sister group to the Metazoa, with Bilateria, Protostomia, Lophotrochozoa, and Ecdysozoa each forming well-supported monophyletic groups. These rooted trees provide strong evidence that the sponge species all belong together in a monophyletic group (Porifera) (bootstrap support = 90% and 96% with outgroup 1 and outgroup 2, respectively). The branch leading to the Porifera is short (Figure 1), accounting for the difficulty in recovering sponge monophyly in previous molecular analyses. This presumably reflects closely spaced splitting events during the Proterozoic era when the sponge lineages emerged.

Extant sponges are a diverse group sharing a number of common body plan features, notably a system of internal canals and choanocyte chambers through which water flows, and a thin epithelial covering called the pinacoderm (Figure 2A). Although morphological character analyses firmly support the hypothesis that the sponges form a monophyletic group [5, 17], rRNA analyses have repeatedly indicated that they are paraphyletic, with the calcisponges and/or the homoscleromorphs positioned closer to eumetazoans than to the other sponges [4–8]. It is worth noting, however, that sponge monophyly could not be ruled out unequivocally in many of these studies because of poor statistical support [6, 7, 10]. The previously proposed hypothesis of sponge paraphyly had significant implications for understanding the origin of multicellular animals, because it would imply that characters shared by all sponge lineages are ancestral for the Metazoa and that eumetazoans are derived from animals with a sponge-like body plan [4, 5, 8, 9] (Figure 2B).

The significant support for sponge monophyly in the present study allows us to return to the idea that a sponge body plan (notably featuring an aquiferous system with internalized

choanocyte chambers and the pinacoderm) evolved in the stem line of the Porifera (Figures 2C and 3). The specialized collar apparatus of sponge choanocytes has often been assumed to be an ancient feature shared with choanoflagellates, based on phenotypic similarity [16]. However, many ultrastructural details of choanoflagellate and choanocyte cells are different, such as the length and spacing of the microvilli and the organization of the microtubule cytoskeleton. Their functional properties also differ, with the microvilli of choanoflagellates but not of choanocytes being contractile. Their similarity might thus represent convergence, with choanocytes being a synapomorphy (shared derived character) of Porifera. It is clear in any case that, rather than reflecting the ancestral animal form, adult sponges are better considered as highly specialized organisms, possibly having acquired a sedentary life style from a hypothetical pelagic ancestor. Notably, the absence of obvious symmetry in many adult sponges fuelled the popular idea that the last common metazoan ancestor lacked defined axial organization [18, 19]. In fact the adult bodies of hexactinellids, calcisponges, homoscleromorphs, and nonbilaterian eumetazoans are characterized by axial symmetry, as is the larval organization of sponges [20], ctenophores, and cnidarians. This suggests that the common ancestor of all animals may have showed symmetry around a single polarity axis [21], and thus that the asymmetry of the adult body in most demosponges and in *Trichoplax* is likely to be derived rather than ancestral (Figure 3).

#### Lessons from Relationships within the Porifera

In line with some previously published phylogenies (e.g., [6, 7, 11]), our analysis placed hexactinellids and demosponges together to form the Silicea Gray, 1867 [22] sensu stricto (with maximal bootstrap support in all analyses) characterized by siliceous spicules organized around a well-defined proteic axial filament [23] and by a particular class of membrane phospholipids known as demosponginic acids [24]. Concerning the enigmatic Homoscleromorpha, our analyses clearly excluded




Figure 3. Changes Affecting Important Body Plan Characters Traced onto the Topology Obtained from Our Molecular Analyses

Key to character changes: 1, acquisition of multicellularity and of a symmetrical body with a single axis of symmetry and polarity; 2, acquisition of the poriferan aquiferous system and of the pinacocytes; 3, acquisition of a well-developed basement membrane supporting epithelia (by convergence in the homoscleromorph sponges and in a cnidarian-ctenophore-bilaterian ancestor); 4, acquisition of siliceous spicules (by convergence in some homoscleromorph sponges and in a hexactinellid + demosponge ancestor, or independently in the hexactinellids and within the demosponges); 5, loss of body symmetry (by convergence in the stem-line of demosponges or within them, and in placozoans); 6, acquisition of gland cells in epithelia [17]; 7, acquisition of the neuro-sensory system, of the muscle cells, and of the digestive system; 8, acquisition of the mesoderm. Homology between the mesoderm of bilaterians, ctenophores, and some cnidarians is debatable; an alternative possibility being convergence of mesoderm-like germ layers between these three taxa; 9, acquisition of bilateral symmetry (by convergence in the Bilateria and in the cnidarian stem-line or within them in the Anthozoa). Parsimony optimization by Mesquite.

them from the demosponges and favored a sister group relationship to the Calcispongiae (with highest support of 91% [BS] obtained in the analysis with outgroup 2), in line with results from 18S rRNA analyses [25, 26] but in conflict with traditional classification schemes (see [27]). The siliceous spicules without defined axial filament found in some Homoscleromorpha [23] thus might have evolved independently from those of hexactinellids and demosponges (Figure 3). In addition, homology of siliceous spicules between the latter two taxa is uncertain because they are absent in the Dictyoceratida, represented here by *Carteriospongia foliascens*, the earliest-branching Demospongiae taxon in our phylogeny (Figure 1) (see [25, 28]). Whether the thick basi-epithelial basement membrane of homoscleromorph larvae and adults, which shares homologous biochemical components with eumetazoan basement membranes [29, 30], was inherited from a common metazoan ancestor and subsequently reduced or lost in most sponges and in *Trichoplax*, or acquired independently in homoscleromorphs and eumetazoans, cannot be decided from our analyses (Figure 3).

### The Coelenterata Clade Revived

A recent phylogenomic analysis suggested that the ctenophores, a phylum of marine, mostly planktonic and gelatinous animals, diverged earlier than sponges [3]. This highly unorthodox hypothesis would see the dismantling of the clade Eumetazoa (ctenophores, cnidarians, and bilaterians), despite their sharing of many key characteristics such as nerve and muscle cells and a differentiated digestive system (absent in

sponges and in *Trichoplax*). Polyphyly of eumetazoans would thus imply several independent acquisitions of these features, or their secondary loss in sponges and/or placozoans [31]. Our rooted analyses are not consistent with the basal position of ctenophores, but rather suggest the existence of a Coelenterata [32] (Ctenophora + Cnidaria) clade, placed within a monophyletic Eumetazoa (Figure 1). A recent study [11] also obtained the coelenterate grouping, but with low bootstrap support, and within a heterodox scheme of eumetazoan polyphyly. Historically, the coelenterate grouping [32] was based on certain anatomical resemblances between ctenophores and the cnidarian medusae (e.g., gelatinous body, tentacles, and “radial” symmetry) that were later considered convergences [33]. In fact, the complex body plan of ctenophores (with eight longitudinal rows of ciliated “comb rows,” a ramified endodermal gastro-vascular system, a complex sensory apparatus located at the aboral pole, and a prevalence of biradial symmetry [19]) differs markedly from that of the cnidarians. Apart from some common embryological features (central yolk and similar unipolar cleavages; animal pole corresponding to adult mouth), there are no clear-cut morpho-anatomical synapomorphies supporting the Coelenterata.

The very long branch leading to the ctenophores (see Figure 1) makes their position prone to perturbation by the long-branch attraction (LBA) artifact [34]. The basal position of ctenophores suggested by Dunn et al. [3] might thus have resulted from attraction of the ctenophores by the distant outgroup taxa used to root the tree. This problem was alleviated in the present study by more comprehensive species sampling

and by the use of the CAT model. That ctenophores are indeed attracted by distant outgroups is empirically demonstrated in our analyses by the observed increase in branch support for Coelenterata and Eumetazoa after partial or total removal of outgroup taxa (Figure 1). Thus, when distant outgroups (notably fungi) were used (as in [3]) (outgroup 1), the Coelenterata were moderately supported (BS = 78%) and the Eumetazoa were poorly supported (BS = 62%). With choanoflagellates as the only outgroup (outgroup 2), support for Coelenterata and Eumetazoa increased remarkably (BS = 93% and BS = 90%, respectively). Even higher support for the coelenterates was obtained by unrooted analysis (BS = 99%). We further checked that the position of ctenophores was not due to artificial attraction by the long branch leading to medusozoan cnidarians (Hydrozoa + Scyphozoa) (see Figure 1), by an analysis excluding these species (Figure S4): ctenophores still grouped with anthozoan cnidarians (a short branch), with high support (BS = 91 %).

Our results not only suggest that ctenophores are the sister group to cnidarians but also that eumetazoans are monophyletic, implying single acquisition during animal evolution of nerve and muscle cells and/or the digestive system, in line with conventional ideas. These findings are at odds with the schemes of eumetazoan polyphyly proposed in two other recent phylogenomic studies [3, 11], both of which used more limited taxonomic sampling of nonbilaterian metazoans and more phylogenetically distant outgroups. It is clearly premature to make a final conclusion on basal metazoan relationships, because not all our analyses yielded significant statistical support values, and the influence of outgroup taxon sampling on tree topology might indicate that there is conflict in the data. As additional data from more nonbilaterian species become available, the remaining doubts should finally be resolved. It should be noted that the position of the placozoan *Trichoplax* with respect to sponges and eumetazoans remains poorly supported in our analyses (Figure 1) and that recent investigations focused on placozoan relationships [11, 35] provided contradictory results, leaving this question unresolved.

### Body Plan Evolution among the Eumetazoans

The proposed restoration of the Coelenterata implies that cnidarians and ctenophores are phylogenetically equally related to the bilaterians and has implications with respect to the origin of mesoderm and of bilateral symmetry. These body plan features have been classically thought to be evolutionary innovations of the Bilateria, but their origin has been suggested to date back to the common cnidarian-bilaterian ancestor from recent developmental gene evidence [36–38]. The mesoderm-like muscle cell lineage of ctenophores [37] might be homologous with the mesoderm of the Bilateria and with mesoderm-like derivatives previously identified in cnidarians [37, 39]. Concerning symmetry, parsimony optimization favors an independent evolution of anatomical bilaterality in the bilaterians and in anthozoan cnidarians (Figure 3), but the significance of the biradial anatomy of the ctenophores [21] remains to be evaluated, for instance through the study of the developmental regulatory genes unilaterally expressed in cnidarians and in the bilaterians [38].

Our new proposal of basal metazoan relationships provides a stimulating framework for furthering our understanding of early metazoan evolution. It suggests that several key features of metazoan body plans were affected by events of convergence or reversion (Figure 3), contrasting with the traditional

conception of metazoan evolution dominated by a gradual increase in morphological complexity. It should motivate detailed exploration of many aspects of character transformations during evolution, development, and metamorphosis, as well as the relationships of larval to adult traits.

### Experimental Procedures

#### EST Sequencing

Fresh samples of *Sycon raphanus*, *Oscarella lobularis*, and *Oopsacas minuta* were collected in the Mediterranean near Marseille (France). *Ephydatia muelleri* gemmules from Belgium were incubated in the lab until production of young adult sponges. Samples of *Heterochone calyx* were collected in British Columbia (Canada) and re-aggregated tissue was used as starting material. *Carterospongia foliascens* was collected at Lizard Island (Great Barrier Reef, Australia) and *Leucetta chagosensis* at North Stradbroke Island (Australia). *Pleurobrachia pileus* adults were collected in Villefranche-sur-Mer (France). For *Clytia hemisphaerica*, the starting material was a strain cultured at the Marine Station in Villefranche-sur-Mer. Frozen samples, RNA Later (QIAGEN)-preserved, or extracted total RNA (depending on the species) were sent to Genome Express (*O. minuta*), RZPD (*S. raphanus*, *O. lobularis*, *E. muelleri*), Express Genomics (*P. pileus* and *C. hemisphaerica*), and the Max Planck Institute for Molecular Genetics in Berlin (Germany) (*H. calyx*, *C. foliascens*, *L. chagosensis*) for cDNA library construction. ESTs were sequenced at the Max Planck Institute for Molecular Genetics (Berlin, Germany) (*H. calyx*, *C. foliascens*, *L. chagosensis*) or at the Genoscope (Evry, France) (all other species). Numbers of sequenced ESTs were approximately 2,000 (*O. minuta*, *E. muelleri*, *S. raphanus*, *O. lobularis*), 4,000 (*H. calyx*, *C. foliascens*, *L. chagosensis*), 30,000 (*P. pileus*), and 90,000 (*C. hemisphaerica*). All these newly sequenced EST collections are publicly available in dbEST/GenBank (<http://www.ncbi.nlm.nih.gov/dbEST/>). The alignment used for phylogenetic analyses is provided as Supplemental Data.

#### Data Assembly

We built upon phylogenomic data sets previously assembled [13, 40]. These alignments were updated, via the protocol described in [41], with the addition of newly generated sequences, and of sequences publicly available from the Trace Archive (<http://www.ncbi.nlm.nih.gov/Traces/>) and the EST Database (<http://www.ncbi.nlm.nih.gov/dbEST/>) of GenBank at the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). In addition, 23 new genes sampled for at least two main poriferan clades were added. All these genes are likely to be orthologs because they are in single-copy in most of the opisthokonts, few recent duplications being observed mostly in vertebrates and *Drosophila*. To further evaluate the possibility of hidden paralogy, we inferred single-gene phylogenies and looked for any strongly supported conflict with the super-matrix tree according to protocol described in [42]. At a bootstrap threshold of 70%, conflicts were observed for only 6.5% of the testable bipartitions, less than the expected error rate. None of these conflicts could be easily explained by hidden paralogy (see details on these analyses in the Supplemental Experimental Procedures).

As previously demonstrated [13], taxon sampling has a major impact in phylogenomic studies. In addition to the nine sponges, nine cnidarians, three ctenophores, and one placozoan available, we therefore selected 22 slowly evolving representative taxa among available Bilateria (based on previous branch length comparison). To reduce the potential impact of long-branch attraction (LBA) [34], we also incorporated all available ichthyosporeans and choanoflagellates (taxa hypothesized to be the closest unicellular relatives of Metazoa) to break the long-branch leading to the distantly related fungal outgroup (for which only the slow-evolving chytridiomycetes and zygomycetes were used).

Ambiguously aligned regions were removed with Gblocks [43]. Sequence selection and concatenation were performed with SCAFoS [44]. To reduce the amount of missing data in the final alignment, we discarded under-sampled genes. Only genes sampled for at least two-thirds of the species (36 out of 55) were retained. The resulting gene selection (128 genes) yielded an alignment of 30,257 unambiguously aligned positions. For all but two genes, the four major diploblast lineages (Porifera, Cnidaria, Ctenophora, and Placozoa) were represented by at least one species; at least three of the main poriferan clades (Demospongiae, Hexactinellida, Homoscleromorpha, Calcispongia) were represented for 65% of the genes.

### Phylogenetic Analyses

To analyze our phylogenomic data set, we used the CAT model [14] with PhyloBayes version 2.3, which has been shown in some contexts to be less prone to LBA artifacts than other models [13, 15]. We performed statistical comparisons of the CAT model with the GTR model (of which other matrix-based models—WAG, JTT, or LG—are special cases) by using cross-validation tests as described in [40]), based on the topology of Figure 1. Ten replicates were run: 9/10 for the learning set and 1/10 for the test set. MCMC were run for 3000 (1500) cycles for the CAT (GTR) model, 1500 (100) being discarded as burn-in. The CAT model was found to have a much better statistical fit than did GTR (a likelihood score of  $3033.36 \pm 123.824$  in favor of CAT). For the plain posterior estimation, two independent chains were run for a total number of 15,000 cycles (corresponding to ~1,200,000 generations) saving every ten cycles and discarding the first 5,000 cycles (burn-in). Each cycle consists in a complicated series of updates of all components of the parameter vector, including an average of 20 topological updates. The posterior consensus tree was obtained by pooling the tree lists of four independent runs. For each node, we compared the posterior probabilities inferred from two independent chains. The maximum difference we observed was 0.11 and 0.2 for the trees with 55 and 44 species, respectively; the average difference being 0.0017 and 0.0011. The differences corresponded to nodes poorly supported by bootstrap values. Posterior probabilities were always identical for nodes having a bootstrap support >90%. Bootstrap percentages were obtained by running 100 independent pseudoreplicates, for 10,000 cycles (900,000 generations) each (because of CPU limitations). Bibliographic references for phylogeny softwares are provided in the Supplemental Experimental Procedures.

### Supplemental Data

Supplemental Data include Supplemental Experimental Procedures, four figures, three tables, and the alignment and can be found with this article online at [http://www.curent-biology.com/supplemental/S0960-9822\(09\)00805-7](http://www.curent-biology.com/supplemental/S0960-9822(09)00805-7).

### Acknowledgments

We thank Muriel Jager for technical help and Ronald Jenner for advice and discussion. We are grateful to Evelyn Richelle-Maurer for providing the *Ephydatia* gemmules, to Thierry Pérez for collecting *Oopsacas minuta* and *Oscarella lobularis* specimens, and to Jean-Yves Carval for collecting *Pleurobrachia pileus* specimens. H.P. acknowledges Génome Québec, the Canadian Research Chair, and the Université de Montréal for financial support and the Réseau Québécois de Calcul de Haute Performance for computational resources. M.M. acknowledges the French Ministry of Research (“ACI jeunes chercheurs” and ANR NT\_NV\_52 Genocnidaire), the Consortium National de Recherche en Génomique, the Genoscope, and the GIS Institut de la Génomique Marine for financial support. S.L. acknowledges funding from NSERC (for the collection and processing of *Heterochone*). G.W. and B.M. acknowledge financial support through the DFG (German Research Foundation) Priority Program SPP1174 “Deep Metazoan Phylogeny” (Project Wo896/6-1,2), M. Kube and his team at the Max-Planck Institute for Molecular Genetics (Berlin, Germany) for library construction and EST sequencing, as well as I. Ebersberger and his team at the Center for Integrative Bioinformatics (Vienna, Austria) for their bioinformatic processing. We also would like to thank Bernard M. Degnan (The University of Queensland, Brisbane, Australia) for providing critical resources that supported cDNA library construction of *Leucetta chagosensis*.

Received: November 13, 2008

Revised: February 22, 2009

Accepted: February 25, 2009

Published online: April 2, 2009

### References

- Philippe, H., and Telford, M. (2006). Large-scale sequencing and the new animal phylogeny. *Trends Ecol. Evol.* 21, 614–620.
- Philippe, H., Lartillot, N., and Brinkmann, H. (2005). Multigene analyses of bilaterian animals corroborate the monophyly of Ecdysozoa, Lophotrochozoa, and Protostomia. *Mol. Biol. Evol.* 22, 1246–1253.
- Dunn, C.W., Hejnol, A., Matus, D.Q., Pang, K., Browne, W.E., Smith, S.A., Seaver, E., Rouse, G.W., Obst, M., Edgecombe, G.D., et al. (2008). Broad phylogenetic sampling improves the resolution of the animal tree of life. *Nature* 452, 745–749.
- Borchiellini, C., Manuel, M., Alivon, E., Boury-Esnault, N., Vacelet, J., and Le Parco, Y. (2001). Sponge paraphyly and the origin of Metazoa. *J. Evol. Biol.* 14, 171–179.
- Peterson, K.J., and Eernisse, D.J. (2001). Animal phylogeny and the ancestry of bilaterians: inferences from morphology and 18S rDNA gene sequences. *Evol. Dev.* 3, 170–205.
- Medina, M., Collins, A.G., Silberman, J.D., and Sogin, M.L. (2001). Evaluating hypotheses of basal animal phylogeny using complete sequences of large and small subunit rRNA. *Proc. Natl. Acad. Sci. USA* 98, 9707–9712.
- Manuel, M., Borchiellini, C., Alivon, E., Le Parco, Y., Vacelet, J., and Boury-Esnault, N. (2003). Phylogeny and evolution of calcareous sponges: Monophyly of Calcinea and Calcaronea, high level of morphological homoplasy, and the primitive nature of axial symmetry. *Syst. Biol.* 52, 311–333.
- Sperling, E.A., Pisani, D., and Peterson, K.J. (2007). Poriferan paraphyly and its implications for Precambrian palaeobiology. In *The Rise and Fall of Ediacaran Biota*, P. Vickers-Rich and P. Komarower, eds. (London: Geological Society Special Publications), pp. 355–368.
- Nielsen, C. (2008). Six major steps in animal evolution: Are we derived sponge larvae? *Evol. Dev.* 10, 241–257.
- Rokas, A., Krüger, D., and Carroll, S.B. (2005). Animal evolution and the molecular signature of radiations compressed in time. *Science* 310, 1933–1938.
- Schierwater, B., Eitel, M., Jakob, W., Osigus, H.-J., Hadrys, H., Dellaporta, S.L., Kolokotronis, S.-O., and DeSalle, R. (2009). Concatenated analysis sheds light on early metazoan evolution and fuels a modern “Urmetzooon” hypothesis. *PLoS Biol.* 7, e1000020.
- Hedtke, S.M., Townsend, T.M., and Hillis, D.M. (2006). Resolution of phylogenetic conflict in large data sets by increased taxon sampling. *Syst. Biol.* 55, 522–529.
- Baurain, D., Brinkmann, H., and Philippe, H. (2007). Lack of resolution in the animal phylogeny: closely spaced cladogeneses or undetected systematic errors? *Mol. Biol. Evol.* 24, 6–9.
- Lartillot, N., and Philippe, H. (2004). A Bayesian mixture model for across-site heterogeneities in the amino-acid replacement process. *Mol. Biol. Evol.* 21, 1095–1109.
- Lartillot, N., Brinkmann, H., and Philippe, H. (2007). Suppression of long-branch attraction artefacts in the animal phylogeny using a site-heterogeneous model. *BMC Evol. Biol.* 7, S4.
- King, N., Westbrook, M.J., Young, S.L., Kuo, A., Abedin, M., Chapman, J., Faircough, S., Hellsten, U., Isogai, Y., Letunic, I., et al. (2008). The genome of the choanoflagellate *Monosiga brevicollis* and the origin of metazoans. *Nature* 451, 783–788.
- Ax, P. (1996). *Multicellular animals: A new approach to the phylogenetic order in nature* (Berlin: Springer-Verlag).
- Hyman, L.H. (1940). *The Invertebrates*, Volume 1–6 (New York: McGraw-Hill).
- Brusca, R.C., and Brusca, G.J. (2003). *Invertebrates*, Second Edition (Sunderland, MA: Sinauer).
- Adamska, M., Degnan, S.M., Green, K.M., Adamski, M., Craigie, A., Larroux, C., and Degnan, B.M. (2007). Wnt and TGF- $\beta$  expression in the sponge *Amphimedon queenslandica* and the origin of metazoan embryonic patterning. *PLoS One* 10, e1031.
- Manuel, M. (2009). Early evolution of symmetry and polarity in metazoan body plans. *C. R. Biol.* 332, 184–209.
- Gray, J.E. (1867). Notes on the arrangement of sponges, with the description of some new genera. *Proc. Zool. Soc. Lond.* 1867, 492–558.
- Uriz, M.-J., Turon, X., Becerro, M.A., and Agell, G. (2003). Siliceous spicules and skeleton frameworks in sponges: Origin, diversity, ultrastructural patterns, and biological functions. *Microsc. Res. Tech.* 62, 279–299.
- Thiel, V., Blumenberg, M., Hefter, J., Pape, T., Pomponi, S., Reed, J., Reitner, J., Wörheide, G., and Michaelis, W. (2002). A chemical view of the most ancient metazoa—biomarker chemotaxonomy of hexactinellid sponges. *Naturwissenschaften* 89, 60–66.
- Borchiellini, C., Chombard, C., Manuel, M., Alivon, E., Vacelet, J., and Boury-Esnault, N. (2004). Molecular phylogeny of Demospongiae: implications for classification and scenarios of character evolution. *Mol. Phylogenet. Evol.* 32, 823–837.

26. Dohrmann, M., Janussen, D., Reitner, J., Collins, A.G., and Wörheide, G. (2008). Phylogeny and evolution of glass sponges (Porifera, Hexactinellida). *Syst. Biol.* 57, 388–405.
27. Hooper, J.N.A., and van Soest, R.W.M., eds. (2002). *Systema Porifera. A Guide to the Classification of Sponges* (New York: Kluwer Academic/Plenum Publishers).
28. Wang, X., and Lavrov, D.V. (2008). Seventeen new complete mtDNA sequences reveal extensive mitochondrial genome evolution within the Demospongiae. *PLoS ONE* 3, e2723.
29. Boute, N., Exposito, J.Y., Boury-Esnault, N., Vacelet, J., Noro, N., Miyazaki, K., Yoshigato, K., and Garrone, R. (1996). Type IV collagen in sponges, the missing link in basement membrane ubiquity. *Biol. Cell* 88, 37–44.
30. Boury-Esnault, N., Ereskovsky, A., Bézac, C., and Tokina, D. (2003). Larval development in Homoscleromorpha (Porifera, Demospongiae). *Invertebr. Biol.* 122, 187–202.
31. Miller, D.J., and Ball, E.E. (2008). Animal evolution: *Trichoplax*, trees, and taxonomic turmoil. *Curr. Biol.* 18, R1003–R1005.
32. Leuckart, R. (1848). *Über die Morphologie und die Verwandtschaftsverhältnisse der wirbellosen Tiere* (Braunschweig: Vieweg und Sohn).
33. Harbison, G.R. (1985). On the classification and evolution of the Ctenophora. In *The Origins and Relationships of Lower Invertebrates. Systematics Association Special, Volume 28*, S.C. Morris, J.D. George, R. Gibson, and H.M. Platt, eds. (Oxford: Clarendon Press), pp. 78–100.
34. Felsenstein, J. (1978). Cases in which parsimony or compatibility methods will be positively misleading. *Syst. Biol.* 27, 401–410.
35. Srivastava, M., Begovic, E., Chapman, J., Putnam, N.H., Hellsten, U., Kawashima, T., Kuo, A., Mitros, T., Salamov, A., Carpenter, M.L., et al. (2008). The *Trichoplax* genome and the nature of placozoans. *Nature* 454, 955–960.
36. Finnerty, J.R., Pang, K., Burton, P., Paulson, D., and Martindale, M.Q. (2004). Origins of bilateral symmetry: *Hox* and *Dpp* expression in a sea anemone. *Science* 304, 1335–1337.
37. Martindale, M.Q. (2005). The evolution of metazoan axial properties. *Nat. Rev. Genet.* 6, 917–927.
38. Matus, D.Q., Pang, K., Marlow, H., Dunn, C.W., Thomsen, G.H., and Martindale, M.Q. (2006). Molecular evidence for deep evolutionary roots of bilaterality in animal development. *Proc. Natl. Acad. Sci. USA* 103, 11195–11200.
39. Seipel, K., and Schmid, V. (2006). Mesodermal anatomies in cnidarian polyps and medusae. *Int. J. Dev. Biol.* 50, 589–599.
40. Lartillot, N., and Philippe, H. (2008). Improvement of molecular phylogenetic inference and the phylogeny of Bilateria. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363, 1463–1472.
41. Baptiste, E., Brinkmann, H., Lee, J.A., Moore, D.V., Sensen, C.W., Gordon, P., Durufle, L., Gaasterland, T., Lopez, P., Muller, M., et al. (2002). The analysis of 100 genes supports the grouping of three highly divergent amoebae: *Dictyostelium*, *Entamoeba*, and *Mastigamoeba*. *Proc. Natl. Acad. Sci. USA* 99, 1414–1419.
42. Rodriguez-Ezpeleta, N., Brinkmann, H., Burger, G., Roger, A.J., Gray, M.W., Philippe, H., and Lang, B.F. (2007). Toward resolving the eukaryotic tree: the phylogenetic positions of jakobids and cercozoans. *Curr. Biol.* 17, 1420–1425.
43. Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol. Biol. Evol.* 17, 540–552.
44. Roure, B., Rodriguez-Ezpeleta, N., and Philippe, H. (2007). SCaFoS: A tool for selection, concatenation and fusion of sequences for phylogenomics. *BMC Evol. Biol.* 7 (Suppl 1), S2.